

# Cryptography

## Lecture 20

# Announcements

- HW 7 due 4/26

# Agenda

- Last time:
  - Number theory
  - Hard problems (Factoring, RSA)
- This time:
  - More number theory (cyclic groups)
  - Hard problems (Discrete log and Diffie-Hellman problems)
  - Elliptic Curve groups

# Cyclic Groups

For a finite group  $G$  of order  $m$  and  $g \in G$ , consider:

$$\langle g \rangle = \{g^0, g^1, \dots, g^{m-1}\}$$

$\langle g \rangle$  always forms a cyclic subgroup of  $G$ .

However, it is possible that there are repeats in the above list.

Thus  $\langle g \rangle$  may be a subgroup of order smaller than  $m$ .

If  $\langle g \rangle = G$ , then we say that  $G$  is a **cyclic group** and that  $g$  is a **generator** of  $G$ .

# Examples

Consider  $Z_{13}^*$ :

2 is a generator of  $Z_{13}^*$ :

$2^0$	1
$2^1$	2
$2^2$	4
$2^3$	8
$2^4$	$16 \rightarrow 3$
$2^5$	6
$2^6$	12
$2^7$	$24 \rightarrow 11$
$2^8$	$22 \rightarrow 9$
$2^9$	$18 \rightarrow 5$
$2^{10}$	10
$2^{11}$	$20 \rightarrow 7$
$2^{12}$	$14 \rightarrow 1$

3 is not a generator of  $Z_{13}^*$ :

$3^0$	1
$3^1$	3
$3^2$	9
$3^3$	$27 \rightarrow 1$
$3^4$	3
$3^5$	9
$3^6$	$27 \rightarrow 1$
$3^7$	3
$3^8$	9
$3^9$	$27 \rightarrow 1$
$3^{10}$	3
$3^{11}$	9
$3^{12}$	$27 \rightarrow 1$

# Definitions and Theorems

Definition: Let  $G$  be a finite group and  $g \in G$ . The order of  $g$  is the smallest positive integer  $i$  such that  $g^i = 1$ .

**Ex:** Consider  $Z_{13}^*$ . The order of 2 is 12. The order of 3 is 3.

Proposition 1: Let  $G$  be a finite group and  $g \in G$  an element of order  $i$ . Then for any integer  $x$ , we have  $g^x = g^{x \bmod i}$ .

Proposition 2: Let  $G$  be a finite group and  $g \in G$  an element of order  $i$ . Then  $g^x = g^y$  iff  $x \equiv y \pmod{i}$ .



# More Theorems

Proposition 3: Let  $G$  be a finite group of order  $m$  and  $g \in G$  an element of order  $i$ . Then  $i \mid m$ .

Proof:

- We know by the generalized theorem of last class that  $g^m = 1 = g^0$ .
- By Proposition 2, we have that  $0 \equiv m \pmod{i}$
- By definition of modulus, this means that  $i \mid m$ .

Corollary: if  $G$  is a group of prime order  $p$ , then  $G$  is cyclic and all elements of  $G$  except the identity are generators of  $G$ .

Why does this follow from Proposition 3?

Theorem: If  $p$  is prime then  $Z_p^*$  is a cyclic group of order  $p - 1$ .



# Prime-Order Cyclic Groups

Consider  $Z_p^*$ , where  $p$  is a strong prime.

- Strong prime:  $p = 2q + 1$ , where  $q$  is also prime.
- Recall that  $Z_p^*$  is a cyclic group of order  $p - 1 = 2q$ .

The subgroup of quadratic residues in  $Z_p^*$  is a cyclic group of prime order  $q$ .

# Example of Prime-Order Cyclic Group

Consider  $Z_{11}^*$ .

Note that 11 is a strong prime, since  $11 = 2 \cdot 5 + 1$ .

$g = 2$  is a generator of  $Z_{11}^*$ :

$2^0$	1
$2^1$	2
$2^2$	4
$2^3$	8
$2^4$	16 $\rightarrow$ 5
$2^5$	10
$2^6$	20 $\rightarrow$ 9
$2^7$	18 $\rightarrow$ 7
$2^8$	14 $\rightarrow$ 3
$2^9$	6

The even powers of  $g$  are the “quadratic residues” (i.e. the perfect squares). Exactly half the elements of  $Z_p^*$  are quadratic residues.

Note that the even powers of  $g$  form a cyclic subgroup of order  $\frac{p-1}{2} = q$ .

Verify:

- closure (Multiplication translates into addition in the exponent. Addition of two even numbers mod  $p - 2$  gives an even number mod  $p - 1$ , since for prime  $p > 3$ ,  $p - 1$  is even.)
- Cyclic –any element is a generator. E.g. it is easy to see that all even powers of  $g$  can be generated by  $g^2$ .

# The Discrete Logarithm Problem

The discrete-log experiment  $DLog_{A,G}(n)$

1. Run  $\mathbf{G}(1^n)$  to obtain  $(G, q, g)$  where  $G$  is a cyclic group of order  $q$  (with  $||q|| = n$ ) and  $g$  is a generator of  $G$ .
2. Choose a uniform  $h \in G$
3.  $A$  is given  $G, q, g, h$  and outputs  $x \in \mathbb{Z}_q$
4. The output of the experiment is defined to be 1 if  $g^x = h$  and 0 otherwise.

Definition: We say that the DL problem is hard relative to  $\mathbf{G}$  if for all ppt algorithms  $A$  there exists a negligible function  $neg$  such that

$$\Pr[DLog_{A,G}(n) = 1] \leq neg(n).$$

# The Diffie-Hellman Problems

# The CDH Problem

Given  $(G, q, g)$  and uniform  $h_1 = g^{x_1}, h_2 = g^{x_2}$ ,  
compute  $g^{x_1 \cdot x_2}$ .

# The DDH Problem

We say that the DDH problem is hard relative to  $G$  if for all ppt algorithms  $A$ , there exists a negligible function  $neg$  such that

$$|\Pr[A(G, q, g, g^x, g^y, g^z) = 1] - \Pr[A(G, q, g, g^x, g^y, g^{xy}) = 1]| \leq neg(n).$$



# Relative Hardness of the Assumptions

Breaking DLog  $\rightarrow$  Breaking CDH  $\rightarrow$  Breaking DDH

DDH Assumption  $\rightarrow$  CDH Assumption  $\rightarrow$  DLog Assumption